



## ASSOCIATION OF THYROID HORMONE PROFILE WITH RESUMPTION OF POSTPARTUM OVARIAN ACTIVITY IN DAIRY COWS

M. YARI<sup>1</sup>, M. KHODAEI-MOTLAGH<sup>1</sup>, M. YAHYAEI<sup>1</sup> & E. DIRANDEH<sup>2</sup>

<sup>1</sup>Department of Animal Science, Faculty of Agriculture and Natural Resources, Arak University, Arak, 38156-8-8349 Iran, <sup>2</sup>Department of Animal Science, Sari Agricultural Sciences and Natural Resources University, P.O. Box 578, Sari, Mazandaran, Iran

### Summary

Yari, M., M. Khodaei-Motlagh, M. Yahyaei & E. Dirandeh, 2021. Association of thyroid hormone profile with resumption of postpartum ovarian activity in dairy cows. *Bulg. J. Vet. Med.* (online first).

The objective of this study was to assess the association between thyroid hormone profile and resumption of postpartum ovulations in cattle. Lactating Holstein cows (n=40, milk yield 38.51±1.23 kg/d) were selected and based on the ultrasound results, blood estradiol and progesterone analysis, were divided into two groups. The cows were classified into anovulatory (AO) group if the first ovulation did not occur until 45 days after calving (n=16) and ovulatory group (O) if the first ovulation occurred 45 days or less after calving (n=24). Blood samples were collected from the cows weekly from day 21 to 48 postpartum to evaluate thyroid hormone levels and blood parameters. Results showed that milk yield and glucose concentrations did not differ between the groups. Serum estradiol and progesterone concentrations in ovulatory cows were significantly higher than those in AO cows. Cows in O group had better condition with respect to days to first service (DFS), open days (OD), days to first ovulation (DFO) and conception rate than cows in AO group. Thyroid hormone analysis show that T<sub>4</sub> concentration and T<sub>4</sub>:T<sub>3</sub> ratio in the AO group were significantly higher than those in O group (P=0.001), while there were no statistically significant differences between groups for T<sub>3</sub> concentrations. Also, T<sub>3</sub> and T<sub>4</sub> concentrations and T<sub>4</sub>:T<sub>3</sub> ratio in both groups were not affected by time (P=0.17) and time×group interaction (P=0.25). In addition, no significant difference was found in the non-esterified fatty acids (NEFA) concentration between the two groups. Unlike NEFA, beta-hydroxybutyric acid (BHBA) levels in AO group were significantly higher than those in O group (P=0.01). In conclusion, cows with different ovarian activity postpartum had different thyroid hormone profile. Increased T<sub>4</sub> concentration and T<sub>4</sub>:T<sub>3</sub> ratio were associated with delay in ovulation and resulted in greater day to first service and open days and followed by lower conception rate

**Key words:** dairy cow, ovarian cyclicity, parturition, thyroid hormone

### INTRODUCTION

Early resumption of the postpartum ovarian cycle in dairy cows is closely related

to reproductive performance in the subsequent period. Cows that ovulated before

21 days postpartum showed better reproductive performance than cows that ovulated for the first time between days 21 and 49 (Rostami *et al.*, 2011). Most cows will have a dominant follicle within 10 to 14 days after calving; however, only 40% of these dominant follicles enter the ovulatory stage (Crowe *et al.*, 2014). In previous studies, ovulation of the first dominant follicle after calving increased estradiol pre-ovulation, whereas most cows that had no higher concentrations of estradiol did not ovulate (Beam & Butler, 1997; Butler *et al.*, 2006). A report on English cows between 1975 and 1982 showed that cows with abnormal ovarian cycles during the postpartum period had longer open days, greater insemination times, and lower conception rates at first insemination compared to normal cows (Darwash *et al.*, 1997). After calving, the intense negative energy balance and accompanying metabolic changes were associated with a decrease in fertility in high-producing dairy herds (Steinhoff *et al.*, 2019). Cows with normal ovarian cycle showed luteal activity sooner than abnormal cows and had a shorter time to first ovulation, although between these cows, a greater distance from calving to first signs of estrus was seen (Ghanem *et al.*, 2014; Sina *et al.*, 2018).

The thyroid axis in the body is one of the important axes for pregnancy in cows and metabolic adaptation during lactation. Thyroid hormones have a major effect on pregnancy and lactation (Steinhoff *et al.*, 2019). Also, thyroid hormones are effective in estrus cycle, pregnancy rate, abortion and stillbirth (Silva *et al.*, 2018). The presence of triiodothyronine (T<sub>3</sub>) and thyroxine (T<sub>4</sub>) in the ovarian follicular fluid of farm animals and thyroid hormone receptor in human and porcine oocytes has been established (Mutinati *et al.*, 2013).

Thyroglobulin (TBG) and TSH receptor have been found in bovine luteal cells (Mutinati *et al.*, 2010). On the other hand, the role of T<sub>3</sub> and T<sub>4</sub> in the steroid production in bovine follicular and luteal cells has been demonstrated (Spicer *et al.*, 2001). Thyroid hormones play an important role in fertility, tissue differentiation and foetal growth, act directly on the ovary, uterus and placenta and promote growth and metabolism through specific receptors on the growth and metabolism of these organs (Silva *et al.*, 2018). Changes in thyroid hormone levels in either direction have an impact on fertility (Mintziori *et al.*, 2016).

Disruption of activity of thyroid gland causes problems in the process of germ cell proliferation and differentiation (Nikraves & Jalaliet, 2006). This significant germ cell proliferation reduction is probably due to the disruption of gonadotropins secretion (Krassas & Pontikides, 2004). There are many conflicting data regarding the effects of hyperthyroidism on reproductive status (Krassas & Pontikides, 2004). In mice, hyperthyroidism alters placental morphology, increases its proliferative activity (Freitas *et al.*, 2007) and affects endometrial status (Kong *et al.*, 2015). Thus, deficiency in thyroid activity has many morphological and physiological effects on reproductive performance of humans and animals. The aim of this study was to investigate the relationship between thyroid hormone concentrations and ovulation in high-producing dairy cows in early lactation.

## MATERIALS AND METHODS

The study was conducted at North part of Iran in Mazandaran Province, Sari (longitude °E 53.06 and latitude °N 36.33). During the study, the ambient temperature

and the relative humidity of the region were 11–24 °C and 52–70%, respectively.

#### *Animals*

Multiparous Holstein dairy cows (n=40, a milk yield of 38.51±1.23, mean±SD) were enrolled into the study at day 10 postpartum. Cows averaged 3.5±1.2 (mean±SD) in parity and 3.12±0.25 (mean±SD) in body condition score at calving based on a 1 (thin) to 5 (obese) scale.

The health status of cows was examined by a veterinarian before the project began. Cows were kept alone with proper ventilation and sand bed in the postpartum period. Feed was offered as a total mixed ration (TMR) formulated according to NRC2001 recommendation 4 times daily (7, 11, 15 and 19 hours) and the same diet was fed throughout the experiment. Water was freely available.

#### *Blood sampling and analyses*

Blood samples were taken weekly from the caudal vein of all cows from day 21 to 48 postpartum (9.0 AM, one hour after morning feeding). Blood serum was then separated and stored at –20 °C until analysis.

Plasma concentrations of progesterone and estradiol  $\beta$ 17 were measured using ELISA Reader (STAT FAX 2100). For this purpose, the ELISA Diaplace Made in Canada kit with kit numbers 4810185 and 49101123 were used. Intra- and inter-assay coefficients of variation were <5%.

Thyroid hormone concentrations in blood plasma were measured by ELISA Reader (STAT FAX 2100) using commercial kits (Orion diagnostica, Finland kit for T<sub>3</sub> and the Monobind ELISA kit (USA) for T<sub>4</sub>). Intra- and inter-assay coefficients of variation were <5%. NEFA (FA 115, Randox Laboratories Ltd., Antrim, UK), and BHBA (Abbott Diabetes

Care Ltd., Witney, UK) were determined by enzymatic colorimetric assays using a spectrophotometer (Shimadzu 2100, Kyoto, Japan).

#### *Evaluation of ovarian dynamics*

Ovarian follicular dynamics were monitored by ultrasound (Easy Scan, BCF Technology, and Livingston, UK) equipped with a 7.5-MHz linear rectal transducer twice weekly from day 21 to 48 postpartum to assess follicular dynamics (dominant follicle diameter, corpus luteum, and ovulation). The change in colour and shape of a large follicle from the prognosis was considered to be the dominant ovulatory follicle. One or two large follicles with a number of small follicles were considered as a follicular wave (Kafi *et al.*, 2015; Nazari *et al.*, 2019).

Cows with at least two consecutive blood samples with a progesterone concentration greater than 1 ng/mL were considered cows with luteal activity (Stevenson, 1997).

When estradiol concentrations were greater than 2 pg/mL and the dominant follicle diameter was greater than or equal to 10 mm, it was considered as the time of ovulation. If the dominant follicle diameter was greater than or equal to 15 mm, but estradiol concentrations were less than or equal to 2 pg/mL, cows were included in the anovulatory group (Cheong *et al.*, 2016).

#### *Study groups*

The base of ultrasound, estradiol and progesterone concentrations results (Kafi *et al.*, 2015), the cows were divided into the following experimental groups:

1. Ovulatory group (O): the first ovulation occurring 45 days or less after calving (n=24).

2. Anovulatory group (AO): the first

ovulation did not occur until 45 days after calving (n=16).

*Statistical analysis*

T<sub>3</sub> and T<sub>4</sub> concentrations in blood plasma samples were examined to investigate the relationship between thyroid hormones and different patterns of luteal activity. Data were analysed using PROC Mixed and SAS software version 9/9. Repeated Measure ANOVA was used to compare data. The statistical model of the design was as follows:

$$Y_{ijk} = \mu + T_i + A_{(i)j} + S_k + TS_{jk} + e_{ijk}$$

where: Y<sub>ijk</sub> = dependent variable; μ = total mean; T<sub>i</sub> = treatment effect; A (i) j = random animal effect in treatment; S<sub>k</sub> = sampling time, TS<sub>jk</sub> = treatment\*time interaction and e<sub>ijk</sub> = is the residual error (Kafi *et al.*, 2012).

**RESULTS**

Milk yield of groups (mean±SD) was similar (49.32±0.56 vs. 50.23±0.49 kg/d). Cows in O group had greater blood estradiol (90.92±2.46 vs. 67.56±3.53) and progesterone (24.32±1.23 vs. 12.32±2.01) concentrations compared to AO cows respectively (P<0.01). The postpartum luteal activity significantly affected DFS, OD, DFO and conception rate (Table 1)

so that cows in O group had a better status compared to AO cows with respect to these parameters.


A significant difference was observed in the T<sub>4</sub> concentration between the two groups (P=0.006), as AO cows had higher T<sub>4</sub> concentration (56.31±2.12 nmol/L) than ovulatory cows (42.64±3.31 nmol/L, Fig. 1). Also, in both groups, T<sub>4</sub> concentration was not affected by time (P=0.46) and time×treatment interaction (P=0.92, Fig. 1). The mean T<sub>3</sub> concentration in O and AO cows was 1.9±1.1 nmol/L and 1.96±0.02 nmol/L respectively and there was no significant difference between them (Fig. 1). Similar to T<sub>4</sub>, time (P=0.17) and time×treatment interaction (P=0.25) did not affect T<sub>3</sub> concentration (Fig. 1).

The mean T<sub>4</sub>:T<sub>3</sub> ratio in AO group was significantly higher (P=0.001) than in O group (Fig. 2). T<sub>4</sub>:T<sub>3</sub> ratio increased significantly over time (P=0.04), while this ratio was not affected by time×treatment interaction (P=0.50).

The results of serum energy metabolites showed no statistically significant difference in NEFA concentrations between the groups (Fig. 3) while the mean BHBA concentrations in AO cows (9.34 ± 0.25 mmol/L) exceeded those in cows from the O group (7.5±0.31 mmol/L). BHBA and NEFA concentrations was not affected by time and time×treatment inter-

**Table 1.** Days to first service, open days, day to first ovulation and conception rate in cows with different postpartum luteal activity patterns. Data are presented as mean ± SD

Item	Postpartum luteal activity pattern	
	Ovulatory group (O) (n=24)	Anovulatory group (AO) (n=16)
Days to first service (DFS)	59.41±0.92 <sup>b</sup>	117.32±0.75 <sup>a</sup>
Open days (OD)	102.26±0.72 <sup>b</sup>	167.55±0.85 <sup>a</sup>
Day to first ovulation (DFO)	39.71±0.62 <sup>b</sup>	95.62±0.62 <sup>a</sup>
Conception rate (%)	69.32±0.25 <sup>a</sup>	30.00±0.25 <sup>b</sup>



**Fig. 1.** Serum T<sub>4</sub> and T<sub>3</sub> levels (mean±SD) of high-producing dairy cows in ovulatory (n=24) and anovulatory groups (n=16) at different postpartum time interval (days); \* P<0.05.


action in both groups. Blood glucose concentration was lower in AO compared to O group ( $4.16 \pm 0.37$  vs.  $4.5 \pm 0.32$  mmol/L) but this difference was not significant ( $P > 0.05$ ).

## DISCUSSION


The postpartum luteal activity affected significantly DFS, OD, DFO and conception rate. In line with our results, previous studies showed a significant correlation between serum progesterone profiles with

a greater incidence of irregular patterns of luteal activity (Gautam *et al.*, 2010). Especially, Taylor *et al.* (2003) found a high correlation between low progesterone concentrations, long periods of anestrus postpartum and delays in uterine involution in dairy cows.

The energy and thyroid hormones levels in the PP period are the key factors that affected the early resumption of ovarian cyclicity in dairy cow after parturition (Gy *et al.*, 2002; Teixeira *et al.*, 2017; Fi-



**Fig. 2.** Serum T<sub>4</sub>:T<sub>3</sub> ratio (mean±SD) of high-producing dairy cows in ovulatory (n=24) and anovulatory groups (n=16) at different postpartum time interval (days); \* P<0.05.



**Fig. 3.** Serum BHBA and NEFA concentrations (mean±SD) of high-producing dairy cows in ovulatory (n=24) and anovulatory groups (n=16) at different postpartum time interval (days); \* P<0.05.

ore *et al.*, 2018). Therefore, the assessment of energy and thyroid hormones profiles can be very important for monitoring ovarian activity in postpartum cows. Based on our knowledge, a few studies have been conducted to investigate the relationship between the level of thyroid hormones and resumption of ovarian cyclicity in dairy cows after parturition (Kafi *et al.*, 2012; Teixeira *et al.*, 2017).

In this study, the cows of the ovulatory group produced the first ovulation until 40 days after parturition which was a better result than that seen in the cows of the AO group where the first ovulation did not occur until 45 days after calving. Also, the cows in the O group had significantly lower concentration of T<sub>4</sub> and T<sub>4</sub>:T<sub>3</sub> ratio compared with the cows in the AO group. These results demonstrated a relationship between high concentration of T<sub>4</sub> and de-

layed first ovulation. Serum thyroid hormones were shown to regulate the reproduction hormones concentration (Gy *et al.*, 2002; Fiore *et al.*, 2018; Steinhoff *et al.*, 2019). An inhibitory effect of thyroid hormones on gonadotropin-releasing hormone (GnRH) and LH secretion in ewes at the transition from the breeding season to anestrus was shown (Anderson *et al.*, 2002). Hence, it could be hypothesised that the higher serum concentrations of  $T_4$  in the postpartum period can lead to delayed first ovulation after parturition in high-producing dairy cows, due to disturbance of the GnRH and LH secretion (Kafi *et al.*, 2012). The high concentration of  $T_4$  in AO group may be attributed to one or both of the following causes: increased production of  $T_4$  and decreased  $T_4$  to  $T_3$  conversion (Kafi *et al.*, 2012). However, the second reason is more probable.

The production rate of thyroid hormones in cattle is influenced by a variety of physiological conditions. It has been proven that high producing dairy cows are in negative energy balance (NEB) during the post-partum period (Gueorguiev, 1999; Jorritsma *et al.*, 2003; Gross *et al.*, 2011). Also, the increased thyroid hormones production at that time is unfavourable due to their negative effects on physiological adjustment and milk production (Gueorguiev, 1999; Fiore *et al.*, 2017). Supposingly, in our study, the higher  $T_4$  concentrations in the AO group compared with O group were not associated with its increased production rate. Since there was a negative correlation between thyroid hormones concentrations and milk yield and blood glucose concentration during the early lactation, uniform average milk yield and blood glucose concentrations in two study groups is one of the reasons that show thyroid hormone production rate is equal in two study

groups. The significantly higher concentration of BHBA in the AO group compared with the O group confirmed this hypothesis. It is known that BHBA is one of the main serum indicators of NEB in dairy cows (Reist *et al.*, 2002; Teixeira *et al.*, 2017) with a negative correlation between BHBA and  $T_3$  and  $T_4$  concentrations (Jorritsma *et al.*, 2003; Kafi *et al.*, 2012; Mohebbi-Fani *et al.*, 2019). So, it was expected that  $T_4$  concentration in the AO group was lower than that in O group while this association was not observed in our study. According to Vap & Weisser (2007) report, BHBA concentrations above 1.4 mmol/L are indicating NEB. In this study minimum concentration of BHBA concentration in AO and O groups were 7.9 mmol/L and 6.4 mmol/L respectively, proving that NEB has occurred in both groups. Several studies have reported the negative influence of BHBA on the resumption of ovarian cyclicity. Consistent with our findings, Kafi *et al.* (2012) showed that anovulatory cows had higher BHBA concentration than cows with normal luteal activity. Mohebbi-Fani *et al.* (2019) reported a positive correlation between BHBA concentration and the interval between calving to first heat. Decreased activity of the 5'-deiodinase enzyme in cows with AO was probably the major reason for the higher  $T_4$  concentration in AO group compared with O group. The 5'-deiodinase enzyme converts  $T_4$  into metabolically active  $T_3$  in the peripheral tissues such as mammary glands and liver (Pezzi *et al.*, 2003). It has been shown the 5'-deiodinase activity can be affected by physiological conditions, for instance, Pezzi *et al.* (2003) showed a clear decrease in liver 5'-deiodinase at the peak of lactation in dairy cows. Samanc *et al.* (2010) reported that 5'-deiodinase activity was decreased in cows with se-

vere fatty liver. Additionally, there was a negative correlation between  $T_3$  concentration and degree of fat accumulation in liver. In our study, the comparison of BHBA between the two groups indicated that cows in the AO group presumably were more susceptible to fatty liver compared with cows in the O group. Therefore, we suggested 5'-deiodinase activity was decreased in AO group that led to a decreased  $T_4$  to  $T_3$  conversion rate of and thus, to high  $T_4$  serum concentration and high  $T_4:T_3$  ratio. Decreased conversion of  $T_4$  to  $T_3$  can affect ovary functions. In the AO group  $T_4:T_3$  ratio showed a significant trend towards increase over time, which result probably shows that as lactation proceeds, 5'-deiodinase activity in liver was decreased. There are studies consistent with our results. Spicer *et al.* (2001) showed a highly significant association of  $T_3$  levels and resumption of ovulation in dairy cows. Also, Pezzi *et al.* (2003) indicated decreased 5'-deiodinase activity at the peak of lactation in dairy cows. The study of Khal *et al.* (1987) confirmed that as lactation proceeded, there was a decrease in liver 5'-deiodinase activity.

Despite the observation that  $T_4$  concentrations were significantly different between the two groups, there was no relevant difference in  $T_3$  concentrations. Consistent with our finding, Kafi *et al.* (2012) showed no significant difference in  $T_3$  concentration between cows with normal and abnormal patterns of luteal activity, whereas  $T_4$  concentration in abnormal cows was significantly higher. Also, it is known that significant changes in peripheral tissue conversion of  $T_4$  to  $T_3$  may occur without a visible change in serum  $T_3$  level (Capuco *et al.*, 1989).

## CONCLUSION

In conclusion, thyroid hormone profiles in dairy cows were associated with ovarian activity postpartum. Increased  $T_4$  concentration and  $T_4:T_3$  ratio were associated with delay in ovulation and resulted in greater day to first service and open days and followed that lower conception rate.

## REFERENCES

- Anderson, G. M., J. M. Connors, S. L. Hardy, M. Valent & R. L. Goodman, 2002. Thyroid hormones mediate steroid-independent seasonal changes in luteinizing hormone pulsatility in the ewe. *Biology of Reproduction*, **66**, 701–706.
- Beam, S. W. & W. Butler, 1997. Energy balance and ovarian follicle development prior to the first ovulation postpartum in dairy cows receiving three levels of dietary fat. *Biology of Reproduction*, **56**, 133–142.
- Butler, S. T., S. H. Pelton & W. Butler, 2006. Energy balance, metabolic status, and the first postpartum ovarian follicle wave in cows administered propylene glycol. *Journal of Dairy Science*, **89**, 2938–2951.
- Capuco, A., J. Keys & J. Smith, 1989. Somatotrophin increases thyroxine-5'-monodeiodinase activity in lactating mammary tissue of the cow. *Journal of Endocrinology*, **121**, 205–211.
- Cheong, S. H., O. G. S. Filho, V. A. Absalón-Medina, S. H. Pelton, W. R. Butler & R. O. Gilbert, 2016. Metabolic and endocrine differences between dairy cows that do or do not ovulate first postpartum dominant follicles. *Biology of Reproduction*, **94**, 1–19.
- Crowe, M., M. Diskin & E. Williams, 2014. Parturition to resumption of ovarian cyclicity: comparative aspects of beef and dairy cows. *Animal*, **8**, 40–53.
- Darwash, A., G. Lamming & J. Woolliams, 1997. Estimation of genetic variation in the interval from calving to postpartum

- ovulation of dairy cows. *Journal of Dairy Science*, **80**, 1227–1234.
- Fiore, E., F. Arfuso, M. Giancesella, D. Vecchio, M. Morgante, E. Mazzotta, T. Badon, P. Rossi, S. Bedin & G. Piccione, 2018. Metabolic and hormonal adaptation in *Bubalus bubalis* around calving and early lactation. *PLoS One*, **13**, 1–13.
- Fiore, E., S. Giambelluca, M. Morgante, G. Piccione, I. Vazzana, B. Contiero, T. Orefice, F. Arfuso & M. Giancesella, 2017. Changes in thyroid hormones levels and metabolism in dairy cows around calving. *Acta Veterinaria*, **67**, 318–330.
- Freitas, E., E. Leite, C. Souza, N. Ocarino, E. Ferreira, G. Cassali, M. Gomes & R. Serakides, 2007. Histomorphometry and expression of Cdc47 and caspase-3 in hyperthyroid rat uteri and placentas during gestation and postpartum associated with fetal development. *Reproduction, Fertility and Development*, **19**, 498–509.
- Gautam, G., T. Nakao, K. Yamada & C. Yoshida, C, 2010. Defining delayed resumption of ovarian activity postpartum and its impact on subsequent reproductive performance in Holstein cows. *Theriogenology*, **73**, 180–189.
- Ghanem, M. E., E. Tezuka, B. Devkota, Y. Izaïke & T. Osawa, 2014. Persistence of uterine bacterial infection, and its associations with endometritis and ovarian function in postpartum dairy cows. *Journal of Reproduction and Development*, **61**, 54–60.
- Gross, J., H. A. van Dorland, R. Bruckmaier & F. Schwarz, 2011. Performance and metabolic profile of dairy cows during a lactational and deliberately induced negative energy balance with subsequent realimentation. *Journal of Dairy Science*, **94**, 1820–1830.
- Gueorguiev, I.P., 1999. Thyroxine and triiodothyronine concentrations during lactation in dairy cows. *Annales de zootechnie*, **48**, 477–480.
- Gy, H., M. Kulcsár & P. Rudas, 2002. Clinical endocrinology of thyroid gland function in ruminants: A review of literature. *Veterinari Medicina*, **47**, 191–202.
- Nikravesh M. R., M. Jalali, 2006. the effect of hypothyroidism on spermatogenesis. *Journal of Iranian Anatomical Science*, **4**, 29–35.
- Jorritsma, R., T. Wensing, T. A. Kruip, P. L. Vos & J. P. Noordhuizen, 2003. Metabolic changes in early lactation and impaired reproductive performance in dairy cows. *Veterinary Research*, **34**, 11–26.
- Kafi, M., A. Tamadon & M. Saeb, 2015. The relationship between serum adiponectin and postpartum luteal activity in high-producing dairy cows. *Theriogenology*, **83**, 1264–1271.
- Kafi, M., A. Tamadon, M. Saeb, A. Mirzaei & M. Ansari-Lari, 2012. Relationships between thyroid hormones and serum energy metabolites with different patterns of postpartum luteal activity in high-producing dairy cows. *animal*, **6**, 1253–1260.
- Kong, L., Q. Wei, J. S. Fedail, F. Shi, K. Nagaoka & G. Watanabe, 2015. Effects of thyroid hormones on the antioxidative status in the uterus of young adult rats. *Journal of Reproduction and Development*, **61**, 219–227.
- Krassas, G. E. & N. Pontikides, 2004. Male reproductive function in relation with thyroid alterations. *Best Practice & Research: Clinical endocrinology & metabolism*, **18**, 183–195.
- Mintziori, G., M. Kita, L. Duntas & D. Gouli, 2016. Consequences of hyperthyroidism in male and female fertility: Pathophysiology and current management. *Journal of Endocrinological Investigation*, **39**, 849–853.
- Mohebbi-Fani, M., A. Omidi, A. Mirzaei, S. Nazifi & K. Nowroozi, 2019. A field study on glucose, non-esterified fatty acids, beta-hydroxybutyrate and thyroid hormones in dairy cows during the breeding period in Fars province, Iran. *Iranian journal of veterinary research*, **20**, 55–59.
- Mutinati, M., S. Desantis, A. Rizzo, S. Zizza, G. Ventriglia, M. Pantaleo & R. Sciorsci,

2010. Localization of thyrotropin receptor and thyroglobulin in the bovine corpus luteum. *Animal Reproduction Science*, **118**, 1–6.
- Mutinati, M., A. Rizzo & R. Sciorsci, 2013. Cystic ovarian follicles and thyroid activity in the dairy cow. *Animal Reproduction Science*, **138**, 150–154.
- Nazari, A., E. Dirandeh, Z. Ansari-Pirsaraei & H. Deldar, 2019. Antioxidant levels, copper and zinc concentrations were associated with postpartum luteal activity, pregnancy loss and pregnancy status in Holstein dairy cows. *Theriogenology*, **133**, 97–103.
- Pezzi, C., P. Accorsi, D. Vigo, N. Govoni & R. Gaiani, 2003. 5'-Deiodinase activity and circulating thyronines in lactating cows. *Journal of Dairy Science*, **86**, 152–158.
- Reist, M., D. Erdin, D. Von Euw, K. Tschuemperlin, H. Leuenberger, Y. Chilliard, H. Hammon, C. Morel, C. Philipona & Y. Zbinden, 2002. Estimation of energy balance at the individual and herd level using blood and milk traits in high-yielding dairy cows. *Journal of Dairy Science*, **85**, 3314–3327.
- Rostami, B., A. Niasari-Naslaji, M. Vojgani, D. Nikjou, H. Amanlou & A. Gerami, 2011. Effect of eCG on early resumption of ovarian activity in postpartum dairy cows. *Animal Reproduction Science*, **128**, 100–106.
- Šamanc, H., V. Stojić, D. Kirovski, M. Jovanović, H. Cernescu & I. Vujanac, 2010. Thyroid hormones concentrations during the mid-dry period: An early indicator of fatty liver in holstein-friesian dairy cows. *Journal of Thyroid Research*, **2010**, 1–6.
- Silva, J. F., N. M. Ocarino & R. Serakides, 2018. Thyroid hormones and female reproduction. *Biology of Reproduction*, **99**, 907–921.
- Sina, M., E. Dirandeh, H. Deldar & B. Shohreh, 2018. Inflammatory status and its relationships with different patterns of postpartum luteal activity and reproductive performance in early lactating Holstein cows. *Theriogenology*, **108**, 262–268.
- Spicer, L., J. Alonso & C. Chamberlain, 2001. Effects of thyroid hormones on bovine granulosa and thecal cell function *in vitro*: Dependence on insulin and gonadotropins. *Journal of Dairy Science*, **84**, 1069–1076.
- Steinhoff, L., K. Jung, M. Meyerholz, J. Heidekorn-Dettmer, M. Hoedemaker & M. Schmicke, 2019. Thyroid hormone profiles and TSH evaluation during early pregnancy and the transition period in dairy cows. *Theriogenology*, **129**, 23–28.
- Stevenson, J. S., 1997. Clinical reproductive physiology of the cow. In *Current therapy in large animal theriogenology* (ed. RS Younquist). WB Saunders, Philadelphia, 257–267.
- Taylor, V. J., D. E. Beever, M. J. Bryant & D. C. Wathes, 2003. Metabolic profiles and progesterone cycles in first lactation dairy cows. *Theriogenology*, **59**, 1661–1677.
- Teixeira, H. C. A., E. A. Barbosa, P. L. G. Souto, A. da Silva Mariante & A. F. Ra, 2017. Postpartum hormone and energy profiles and their influence on the resumption of ovarian cyclicity in Curraleiro Pé-Duro cows. *Theriogenology*, **95**, 133–140.
- Vap, L. M. & M. G. Weiser, 2007. Field chemistry analysis. *Veterinary Clinics of North America: Food Animal Practice*, **23**, 427–442.

Paper received 16.11.2020; accepted for publication 09.02.2021

#### Correspondence:

Associate professor Mahdi Khodaei-Motlagh  
Department of Animal Science,  
Faculty of Agriculture and Natural Science,  
Arak University, Arak 38156-8-8349, Iran  
e-mail: mmotlagh2002@gmail.com