



---

## SCREENING FOR SARCOPENIA IN ELDERLY WOMEN

P. Penev\*

University “Prof. Dr. Assen Zlatarov” - Burgas Burgas, Bulgaria

### ABSTRACT

**PURPOSE:** The knowledge for the disease sarcopenia among the elderly population in Bulgaria is insufficient. Whole-body dual-energy X-ray (DXA) scans, anthropometric and clinical tests are used to diagnose it. Early screening of sarcopenia is an important strategy for the timely detection of this disease. The aim of the current study is to identify sarcopenia in the elderly through SARC-F questionnaire.

**METHODS:** Data were obtained based on a questionnaire conducted by the student nurses in the Faculty of Public Health and Health Care of the University "Prof. Dr. Assen Zlatarov"-Burgas. A five-component SARC-F questionnaire was used to assess muscle strength. A sum equal to or greater than 4 is predictive of sarcopenia. 113 women aged  $\geq 65$  with a mean age of  $74.6 \pm 6.7$  years were included in the study.

**RESULTS:** 37% of the women had a total score corresponding to the risk of sarcopenia. The most common signs were a lack of muscle strength and an increased number of falls. The assessment of SARC-F might facilitate the physicians in the clinical practice to detect sarcopenia in the elderly population.

**CONCLUSIONS:** SARC-F is a reliable method for the assessment of sarcopenia in elderly women.

**Key words:** sarcopenia, muscle strength, screening, SARC-F

### INTRODUCTION

The term, sarcopenia, was firstly used in 1988 to describe the condition with muscle wasting of the older people (1). In the next years, authors defined sarcopenia as appendicular muscle mass  $< 2$  standard deviations (SDs) below the mean of a young reference group assessed with dual energy X-ray absorptiometry (DXA). In previous published studies the prevalence and the severity of sarcopenia were reported to increase significantly with age and sarcopenia was associated with a physical disability (2). Through assessment with bioelectrical impedance analysis (BIA) people with functional impairment showed an estimated lean mass below 2 SDs of the mean (3). Obesity and loss of muscle mass were also attributed to worse outcomes. The term

‘sarcopenic obesity’ became popular (4). It was recognized that there are numerous causes of age-related sarcopenia, including loss of motor units innervating muscle, systemic inflammation, oxidative stress, and decline in anabolic hormones and decrease in the physical activity (5, 6).

### PURPOSE

Sarcopenia and osteoporosis are often coexistent. There are different tools for the assessment of osteoporosis and fracture risk to make a decision if the patients should undergo DXA scan (7,8). Several sarcopenia screening tools have been developed to assess this condition. As a pioneer of these tools, the SARC-F has been validated in various studies and translated into different languages (9-11).

The SARC-F has 5 items: strength, assistance in walking, rising from a chair, climbing stairs, and falls, (**Figure 1**)

---

\*Correspondence to: Plamen Valev Penev, University “Prof. Dr. Assen Zlatarov” – Burgas, E-mail address: [osteoblik@abv.bg](mailto:osteoblik@abv.bg), Telephone number: 0887409723

Component	Question	Scoring
Strength	How much difficulty do you have in lifting and carrying 10 pounds?	None = 0 Some = 1 A lot or unable = 2
Assistance in walking	How much difficulty do you have walking across a room?	None = 0 Some = 1 A lot, use aids, or unable = 2
Rise from a chair	How much difficulty do you have transferring from a chair or bed?	None = 0 Some = 1 A lot or unable without help = 2
Climb stairs	How much difficulty do you have climbing a flight of 10 stairs?	None = 0 Some = 1 A lot or unable = 2
Falls	How many times have you fallen in the past year?	None = 0 1–3 falls = 1 4 or more falls = 2

**Figure 1.** SARC-F questionnaire

## METHODS

Data were obtained based on a questionnaire conducted by the student nurses in the Faculty of Public Health and Health Care of the University "Prof. Dr. Assen Zlatarov"-Burgas. Five-component SARC-F questionnaire was used to assess muscle strength, the need of assistance in walking, the ability to rise from a chair and to climb stairs, and the incidence of falls. The score ranges from 0 to 10 points with 0 to 2 points for each component. A sum equal to or greater than 4 is predictive of sarcopenia. 113 women with a mean age of  $74.6 \pm 6.7$  years were included in the study. The inclusion criteria were subjects of age  $\geq 65$  years. Other concomitant diseases were not evaluated. Subjects haven't undergone DXA scans or bioelectrical impedance analysis (BIA). SPSS version 19 was used to do statistical analysis of the data.

## RESULTS


42/113 women (37%) were diagnosed with risk of sarcopenia (the sum of the SARC-F questionnaire was equal or above 4 points). The remain subjects-71/113 women (63%) didn't have risk sarcopenia (the sum of the SARC-F questionnaire was less than 4 points). The age of the subjects correlated with the incidence of sarcopenia. Sarcopenic women were significantly older (77.4 years) than non-sarcopenic (73.2 years),  $p < 0.05$ . The most common signs of the women with sarcopenia were lack of muscle strength and increased number of falls, fig. 2. According to the component "assistance in walking" 43 women (38.1%) showed no difficulty in walking across a room, 61 women had some difficulty (53.9%) and 9 women (8%) had a lot of difficulty in walking across a room or were unable to walk. After assessing the subjects due to the criterion "rise from a chair" we

determined that 38 women (33.6%) were with no difficulty and 42 women (37.2%) were with some difficulty in transferring from a chair or bed. The remaining 33 women (29.2%) were unable to rise from a chair or bed without help. 42 subjects (37.2%) didn't show any difficulties in climbing 10 stairs, 54 subjects (47.8%) had some difficulties in doing this activity and 17 subjects (15%) were unable to climb 10 stairs. 58 women (51.3%) haven't fallen in the past year, 41 women (36,3%) had 1-3 falls and 14 women (12.4%) had 4 or more falls in the past year. 51 subjects (45.1%) had no difficulty in lifting and caring 10 pounds, 44 subjects (38.9%) had some difficulty and 18 subjects (15.9%) had a lot of difficulties or were unable to lift and carry 10 pounds, **Table 1**.

## DISCUSSION

Despite the clinical importance of sarcopenia this disease remains poorly recognized in the clinical practice. Although some studies suggested that cutoff value of 3 of SARC-F reached the optimal Youden's index, sarcopenia is considered to be present if  $SARC-F \geq 4$  (12, 13). SARC-F is a simple method for screening of sarcopenia and could be helpful to detect high-risk patients for further examinations.

The prediction of the adverse outcomes is the most important factor in evaluating the function of screening scores. Tan et al. (14) detected negative health outcomes in patients at high risk of sarcopenia assessed with SARC-F. The results of the current study showed that SARC-F had a good performance in predicting sarcopenia and suggested that early exercise intervention for high-risk individuals detected with SARC-F might reduce the risk of adverse outcomes and improve prognosis.



**Figure 2.** Percent of the women with and without risk of sarcopenia.

**Table 1.** Number and percent of the women according to each separate component of SARC-F

Components		Count of the women	% of the women
assistance in walking	No difficulty	43	38.1%
	Some difficulty	61	53.9%
	A lot of difficulty or unable	9	8.0%
rise from a chair	No difficulty	38	33.6%
	Some difficulty	42	37.2%
	A lot of difficulty or unable	33	29.2%
climb stairs	No difficulty	42	37.2%
	Some difficulty	54	47.8%
	A lot of difficulty or unable	17	15.0%
falls	0	58	51.3%
	1-3	41	36.3%
	4 or more	14	12.4%
strength	No difficulty	51	45.1%
	Some difficulty	44	38.9%
	A lot of difficulty or unable	18	15.9%

Some studies showed very low sensitivity of the SARC-F (15-17). One of them reported that SARC-F had a sensitivity of 35.6% and a specificity of 82.2% in 487 Mexican older adults (18).

In another study which compared SARC-F-3 with SARC-F was found that SARC-F-3 had even lower sensitivity and diagnostic accuracy than SARC-F. Low sensitivity means that subjects who have sarcopenia could be missed. An AUC area of > 0.9 indicates high accuracy, 0.7 to 0.9 indicates moderate accuracy, 0.5 to 0.7-low accuracy and 0.5 indicates chance result (19). In the study of Yang et al., the AUC of SARC-F was 0.894, whereas that of SARC-F-3 was 0.676 and this was the reason why the authors of this study suspected that SARC-F-3 may not be suitable for sarcopenia screening in community-dwelling older adults (20). The incidence of sarcopenia (37%) in our study was higher compared to those found in the studies of Yang et al.-15.9% (19), Smoliner et al.-25% (21) and Coin et al.-20% who assessed community-dwelling people in Italy (22). Masanes et al. evaluated community-dwelling elderly in Barcelona, Spain and found a similar prevalence of sarcopenia (33%) compared to our study (37%).

## CONCLUSIONS

In the current study, we found a relatively high prevalence of sarcopenia compared to previous studies. Sarcopenic women were significantly older than non-sarcopenic.

## REFERENCES

- Rosenberg IH. Sarcopenia: origins and clinical relevance. *J Nutr* 127( Suppl: 990S– 991S, 1997
- Baumgartner RN, Koehler KM, Gallagher D, Romero L, Heymsfield SB, Ross RR, et al. Epidemiology of sarcopenia among the elderly in New Mexico. *Am J Epidemiol*; 147: 755– 763, 1998
- Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatr Soc*; 50: 889– 896, 2002
- Baumgartner RN, Wayne SJ, Waters DL, Janssen I, Gallagher D, Morley JE. Sarcopenic obesity predicts instrumental activities of daily living disability in the elderly. *Obes Res* 12: 1995– 2004, 2004;
- Morley JE, Baumgartner RN, Roubenoff R, Mayer J, Nair KS. Sarcopenia. *J Lab Clin Med* 137: 231– 243, 2001
- Marzetti E, Calvani R, Tosato M, et al. Sarcopenia: an overview. *Aging Clin Exp Res* 29: 11–17, 2017
- Kirilova E, D Tcherkezov, V Gonchev, Z Zheleva OSIRIS index for assessment of the osteoporosis risk in postmenopausal women. National conference Science and Society 2019, Kardjali, RKR print OOD ISSN 1314-3425, 95-99, 2019.
- Kirilov N, Kirilova E. Computer vision for assessment of scoliosis from dual energy X-ray (DXA) image, Second Youth Scientific Session “Biomedicine and Quality of Life” November 4-5, , BMQL-2019-Book-of-abstracts.pdf: p 18, 2019
- Cao L, Chen S, Zou C, et al. A pilot study of the SARC-F scale on screening sarcopenia and physical disability in the Chinese older people. *J Nutr Health Aging*; 18:277–283, 2014
- Parra-Rodriguez L, Szlejf C, Garcia-Gonzalez AI, et al. Cross-cultural adaptation and validation of the Spanish-language version of the SARC-F to assess sarcopenia in Mexican community-dwelling older adults. *J Am Med Dir Assoc*, 17:1142–1146, 2016
- Kim S, Kim M, Won CW. Validation of the Korean version of the SARC-F questionnaire to assess sarcopenia: Korean frailty and aging cohort study. *J Am Med Dir Assoc*, 19: 40–45, 2018
- Malmstrom TK, Morley JE. SARC-F: a simple questionnaire to rapidly diagnose sarcopenia. *J Am Med Dir Assoc*, 14:531– 532, 2013
- Li M, Kong Y, Chen H, Chu A, Song G, Cui Y. Accuracy and prognostic ability of the SARC-F questionnaire and Ishii's score in the screening of sarcopenia in geriatric inpatients. *Braz J Med Biol Res*; 52(9):e8204. doi:10.1590/1414-431X20198204, 2019
- Tan LF, Lim ZY, Choe R, Seetharaman S, Merchant R. Screening for frailty and sarcopenia among older persons in medical outpatient clinics and its associations with healthcare burden. *J Am Med Dir Assoc* 18:583–587, 2017
- Malmstrom TK, Morley JE. SARC-F: a simple questionnaire to rapidly diagnose sarcopenia. *J Am Med Dir Assoc* 14:531– 532, 2013

16. Tanaka S, Kamiya K, Hamazaki N, et al. Utility of SARC-F for assessing physical function in elderly patients with cardiovascular disease. *J Am Med Dir Assoc* 18:176–181, 2017
17. Parra-Rodriguez L, Szejf C, Garcia-Gonzalez AI, et al. Cross-cultural adaptation and validation of the Spanish-language version of the SARC-F to assess sarcopenia in Mexican community-dwelling older adults. *J Am Med Dir Assoc*, 17:1142–1146, 2016
18. Linden A. Measuring diagnostic and predictive accuracy in disease management: an introduction to receiver operating characteristic (ROC) analysis. *J Eval Clin Pract* 12:132–139, 2006
19. Yang M, Hu X, Xie L, et al. SARC-F for sarcopenia screening in community-dwelling older adults: Are 3 items enough?. *Medicine* (Baltimore). 97(30):e11726. doi:10.1097/MD.0000000000011726, 2018
20. Barbosa-Silva TG, Menezes AM, Bielemann RM, et al. Enhancing SARC-F: improving sarcopenia screening in the clinical practice. *J Am Med Dir Assoc* 17:1136–1141, 2016.
21. Smoliner C, Sieber CC, Wirth R. Prevalence of sarcopenia in geriatric hospitalized patients. *J Am Med Dir Assoc*; 15:267–272, 2014.
22. Coin A, Sarti S, Ruggiero E, Giannini S, Pedrazzoni M, Minisola S, et al. Prevalence of sarcopenia based on different diagnostic criteria using DEXA and appendicular skeletal muscle mass reference values in an Italian population aged 20 to 80. *J Am Med Dir Assoc*. ;14:507–512, 2013
23. Masanes F, Culla A, Navarro-Gonzalez M, Navarro-Lopez MC, Sacanella E, Torres B, et al. Prevalence of sarcopenia in healthy community-dwelling elderly in an urban area of Barcelona (Spain) *J Nutr Health Aging*. ;16:184–187, 2012